

# Homework 5

All rings are commutative with unit.

Recall that an ideal  $\mathfrak{q}$  in a ring  $A$  is *primary* if  $xy \in \mathfrak{q}$  implies that either  $x \in \mathfrak{q}$  or  $y^n \in \mathfrak{q}$  for some  $n$ . Equivalently, every zero divisor in  $A/\mathfrak{q}$  is nilpotent.

**Lemma 1.** *If  $\mathfrak{q}$  is primary then  $r(\mathfrak{q})$  is prime.*

*Proof.* Suppose that  $xy \in r(\mathfrak{q})$ . Then  $x^N y^N \in \mathfrak{q}$  for some  $N$ . Thus either  $x^N \in \mathfrak{q}$  or  $(y^N)^n \in \mathfrak{q}$  for some  $n$ . Thus either  $x \in r(\mathfrak{q})$  or  $y \in r(\mathfrak{q})$ .  $\square$

If  $\mathfrak{q}$  is primary and  $r(\mathfrak{q}) = \mathfrak{p}$  then we say that  $\mathfrak{q}$  is  $\mathfrak{p}$ -primary.

**Lemma 2.** *Suppose that  $\mathfrak{q}_1, \mathfrak{q}_2$  are  $\mathfrak{p}$ -primary ideals. Then  $\mathfrak{q}_1 \cap \mathfrak{q}_2$  is also  $\mathfrak{p}$ -primary.*

*Proof.* We have  $r(\mathfrak{q}_1 \cap \mathfrak{q}_2) = r(\mathfrak{q}_1) \cap r(\mathfrak{q}_2) = \mathfrak{p} \cap \mathfrak{p} = \mathfrak{p}$ . To show  $\mathfrak{q}_1 \cap \mathfrak{q}_2$  is  $\mathfrak{p}$ -primary, suppose that  $xy \in \mathfrak{q}_1 \cap \mathfrak{q}_2$  but  $x \notin \mathfrak{q}_1 \cap \mathfrak{q}_2$ . Then either  $x \notin \mathfrak{q}_1$  or  $x \notin \mathfrak{q}_2$ . By symmetry, we may assume that  $x \notin \mathfrak{q}_1$ . Then since  $xy \in \mathfrak{q}_1$  which is  $\mathfrak{p}$ -primary, we have  $y \in \mathfrak{p} = r(\mathfrak{q}_1 \cap \mathfrak{q}_2)$ .  $\square$

**Theorem 3.** *If  $A$  is Noetherian, then every ideal may be expressed as a finite intersection of primary ideals.*

*Proof.* We will prove this in class. See Lang's *Algebra*, Theorem X.3.3 on page 423.  $\square$

Write the ideal  $\mathfrak{a}$  as an intersection of primary ideals  $\mathfrak{q}_i$ :

$$\mathfrak{a} = \bigcap_{i=1}^d \mathfrak{q}_i.$$

By Lemma 2 we may assume that the radicals  $r(\mathfrak{q}_i)$  are all distinct. Moreover, we may obviously discard any  $\mathfrak{q}_i$  if that does not change the intersection. Assuming this, the decomposition is called a (reduced) *primary decomposition*. The decomposition is not unique but has some uniqueness properties. The primary decomposition is discussed in Lang's *Algebra* Chapter 10, but see the book of Atiyah and Macdonald, Chapter 4, for more information. Atiyah and Macdonald is available on-line through the Stanford Libraries.

The notion of a primary ideal is needed in the dimension theory.

The following well-known fact may be useful in doing Problem 1. The set  $\mathfrak{N}$  of nilpotent elements of a ring  $A$  is called the *nilradical*. It is an ideal, the radical of  $(0)$ .

**Proposition 4.** *The nilradical is the intersection of all prime ideals of  $A$ .*

*Proof.* See Lang's *Algebra*, Corollary X.2.2 on page 417.  $\square$

**Problem 1.** Show that if  $r(\mathfrak{a})$  is maximal, then  $A/\mathfrak{a}$  is a local ring with a single prime ideal. The ideal  $\mathfrak{a}$  is primary.

One might hope from Problem 1 that if  $r(\mathfrak{a})$  is only assumed to be prime that  $\mathfrak{a}$  might be primary. However the next problem shows that this is not true.

**Problem 2.** Let  $A$  be the quotient of the polynomial ring  $\mathbb{C}[X, Y, Z]$  by the ideal  $(X^2 - YZ)$  and let  $x, y, z$  be the images of  $X, Y, Z$  in  $A$ . Let  $\mathfrak{p}$  be the ideal  $(x, z)$ . Show that  $\mathfrak{p}$  is prime and that the radical of  $\mathfrak{p}^2$  is  $\mathfrak{p}$ , but that  $\mathfrak{p}^2$  is not primary.

**Problem 3.** Let  $A$  be the polynomial ring  $k[x, y, z]$ , where  $k$  is a field. Let  $\mathfrak{p}_1 = (x, y)$ ,  $\mathfrak{p}_2 = (x, z)$  and  $\mathfrak{m} = (x, y, z)$ . Let  $\mathfrak{a} = \mathfrak{p}_1\mathfrak{p}_2$ . Prove that

$$\mathfrak{a} = \mathfrak{p}_1 \cap \mathfrak{p}_2 \cap \mathfrak{m}^2$$

is a primary decomposition.

The next problem is a strengthening from a fact from last week.

**Problem 4.** Let  $\mathfrak{o}$  be a Dedekind domain with a unique maximal ideal  $\mathfrak{p}$ . Prove that  $R$  is a discrete valuation ring.

**Problem 5.** Let  $A$  be the quotient of the polynomial ring  $k[X, Y]$  by the polynomial  $Y^2 - X^3$  and let  $x, y$  be the images of  $X$  and  $Y$  in  $A$ . Prove that  $A$  is not integrally closed. If  $\mathfrak{m}$  is the maximal ideal generated by  $x$  and  $y$ , determine the dimension of  $\mathfrak{m}^n/\mathfrak{m}^{n+1}$  for all  $n \geq 1$ .

**Problem 6.** Let  $\lambda$  be a function defined on modules over a ring taking values in  $\mathbb{Z} \cup \{\infty\}$  such that if  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  is a short exact sequence then

$$\lambda(M) = \lambda(M') + \lambda(M'').$$

Prove that if  $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow M_4 \rightarrow 0$  is exact then

$$\lambda(M_1) - \lambda(M_2) + \lambda(M_3) - \lambda(M_4) = 0.$$